

Lecture Notes of  
*Seminario Interdisciplinare di Matematica*  
Vol. 9(2010), pp. 111–128.

## The functions of the Wright type in fractional calculus

Francesco MAINARDI, Antonio MURA and Gianni PAGNINI

*Dedicated to Professor Paolo Emilio Ricci on the occasion of his retirement*

**Abstract**<sup>1</sup>. We provide a survey of the high transcendental functions known in the literature as Wright functions. We devote particular attention for two functions of the Wright type, which, in virtue of their role in applications of fractional calculus, we have called auxiliary functions. We also discuss their relevance in probability theory showing their connections with Lévy stable distributions. At the end, we add some historical and bibliographical notes.

### 1. INTRODUCTION

Here we provide a survey of the high transcendental functions related to the Wright special function. Like the functions of the Mittag-Leffler type, the functions of the Wright type are known to play fundamental roles in various applications of the fractional calculus. This is mainly due to the fact that they are interrelated with the Mittag-Leffler functions through Laplace and Fourier transformations. We start providing the definitions in the complex plane for the general Wright function and for two special cases that we call auxiliary functions. Then we devote particular attention to the auxiliary functions in the real field, because they admit a probabilistic interpretation related to the fundamental solutions of certain evolution equations of fractional order. These equations are fundamental to understand phenomena of anomalous diffusion or intermediate between diffusion and wave propagation.

At the end we add some historical and bibliographical notes.

### 2. THE WRIGHT FUNCTION $W_{\lambda,\mu}(z)$

The Wright function, that we denote by  $W_{\lambda,\mu}(z)$  is so named in honour of E. Maitland Wright, the eminent British mathematician, who introduced and investigated this function in a series of notes starting from 1933 in the framework of the

---

<sup>1</sup>Authors' address: F. Mainardi, Università degli Studi di Bologna, Dipartimento di Fisica and INFN, Via Irnerio 46, I-40126 Bologna, Italy; e-mail: [francesco.mainardi@unibo.it](mailto:francesco.mainardi@unibo.it).

A. Mura, CRESME Ricerche S.p.A., Viale Gorizia 25C, I-00199 Roma, Italy;  
e-mail: [anto.mura@gmail.com](mailto:anto.mura@gmail.com).

G. Pagnini, CRS4, Centro Ricerche Studi Superiori e Sviluppo in Sardegna, Polaris Bldg. 1, I-09010 Pula (Cagliari), Italy; e-mail: [pagnini@crs4.it](mailto:pagnini@crs4.it).

*Keywords.* Wright functions, Mittag-Leffler functions, fractional calculus, stable distributions.  
*AMS Subject Classification.* 26A33, 33E12, 33C40, 44A10, 45K05, 60G52.

asymptotic theory of partitions, see Wright [42, 43, 44]. The function is defined by the series representation, convergent in the whole  $z$ -complex plane,

$$(1) \quad W_{\lambda,\mu}(z) := \sum_{n=0}^{\infty} \frac{z^n}{n! \Gamma(\lambda n + \mu)} \quad , \quad \lambda > -1 \quad , \quad \mu \in \mathbb{C} \quad ,$$

so  $W_{\lambda,\mu}(z)$  is an *entire function*. Originally, Wright assumed  $\lambda > 0$ , and, only in 1940, he considered  $-1 < \lambda < 0$ , see Wright [45]. We note that in the handbook of the Bateman Project (Erdélyi et al. [6], Vol. 3, Ch. 18), presumably for a misprint,  $\lambda$  is restricted to be non negative.

**The integral representation.** The *integral representation* reads

$$(2) \quad W_{\lambda,\mu}(z) = \frac{1}{2\pi i} \int_{Ha} e^{\sigma+z\sigma^{-\lambda}} \frac{d\sigma}{\sigma^\mu} \quad , \quad \lambda > -1 \quad , \quad \mu \in \mathbb{C} \quad ,$$

where  $Ha$  denotes the Hankel path, i.e. a loop which starts and ends at  $-\infty$  and encircles the circular disk  $|\zeta| \leq |z|^{1/\alpha}$  in the positive sense:  $-\pi \leq \arg \zeta \leq \pi$  on  $Ha$ . The equivalence of the series and integral representations is easily proved using the Hankel formula for the Gamma function

$$\frac{1}{\Gamma(\zeta)} = \int_{Ha} e^u u^{-\zeta} du \quad , \quad \zeta \in \mathbb{C} \quad ,$$

and performing a term-by-term integration. In fact,

$$\begin{aligned} W_{\lambda,\mu}(z) &= \frac{1}{2\pi i} \int_{Ha} e^{\sigma+z\sigma^{-\lambda}} \frac{d\sigma}{\sigma^\mu} = \frac{1}{2\pi i} \int_{Ha} e^\sigma \left[ \sum_{n=0}^{\infty} \frac{z^n}{n!} \sigma^{-\lambda n} \right] \frac{d\sigma}{\sigma^\mu} = \\ &= \sum_{n=0}^{\infty} \frac{z^n}{n!} \left[ \frac{1}{2\pi i} \int_{Ha} e^\sigma \sigma^{-\lambda n - \mu} d\sigma \right] = \sum_{n=0}^{\infty} \frac{z^n}{n! \Gamma[\lambda n + \mu]} \quad . \end{aligned}$$

It is possible to prove that the Wright function is entire of order  $1/(1+\lambda)$ , hence of exponential type if  $\lambda \geq 0$ . The case  $\lambda = 0$  is trivial since  $W_{0,\mu}(z) = e^z/\Gamma(\mu)$ .

**Asymptotic expansions.** For the detailed asymptotic analysis in the whole complex plane for the Wright functions, the interested reader is referred to Wong and Zhao [40, 41]. These authors have provided asymptotic expansions of the Wright functions of the first ( $\lambda \geq 0$ ) and second ( $-1 < \lambda < 0$ ) kind following a new method for smoothing Stokes' discontinuities.

As a matter of fact, the second kind is the most interesting for us. By setting  $\lambda = -\nu \in (-1, 0)$ , we recall the asymptotic expansion originally obtained by Wright himself, that is valid in a suitable sector about the negative real axis as  $|z| \rightarrow -\infty$ ,

$$(3) \quad \begin{aligned} W_{-\nu,\mu}(z) &= Y^{1/2-\mu} e^{-Y} \left[ \sum_{m=0}^{M-1} A_m Y^{-m} + O(|Y|^{-M}) \right] \quad , \\ Y &= Y(z) = (1-\nu)(-\nu^\nu z)^{1/(1-\nu)} \quad , \end{aligned}$$

where the  $A_m$  are certain real numbers.

**Generalization of the Bessel functions.** The Wright functions turn out to be related to the well-known Bessel functions  $J_\nu$  and  $I_\nu$  for  $\lambda = 1$  and  $\mu = \nu + 1$ . In fact, by using the well-known series definitions for the Bessel functions and the

series definitions (1) for the Wright functions, we easily recognize the identities:

$$\begin{aligned}
 J_\nu(z) &:= \left(\frac{z}{2}\right)^\nu \sum_{n=0}^{\infty} \frac{(-1)^n (z/2)^{2n}}{n! \Gamma(n + \nu + 1)} = \left(\frac{z}{2}\right)^\nu W_{1,\nu+1} \left(-\frac{z^2}{4}\right), \\
 W_{1,\nu+1}(-z) &:= \sum_{n=0}^{\infty} \frac{(-1)^n z^n}{n! \Gamma(n + \nu + 1)} = z^{-\nu/2} J_\nu(2z^{1/2}),
 \end{aligned}
 \tag{4}$$

and

$$\begin{aligned}
 I_\nu(z) &:= \left(\frac{z}{2}\right)^\nu \sum_{n=0}^{\infty} \frac{(z/2)^{2n}}{n! \Gamma(n + \nu + 1)} = \left(\frac{z}{2}\right)^\nu W_{1,\nu+1} \left(\frac{z^2}{4}\right), \\
 W_{1,\nu+1}(z) &:= \sum_{n=0}^{\infty} \frac{z^n}{n! \Gamma(n + \nu + 1)} = z^{-\nu/2} I_\nu(2z^{1/2}).
 \end{aligned}
 \tag{5}$$

As far as the standard Bessel functions  $J_\nu$  are concerned, the following observations are worth noting. We first note that the Wright function  $W_{1,\nu+1}(-z)$  reduces to the entire function  $C_\nu(z)$  known as *Bessel-Clifford function*. Then, in view of the first equation in (4) some authors refer to the Wright function as the *Wright generalized Bessel function* (misnamed also as the *Bessel-Maitland function*) and introduce the notation for  $\lambda \geq 0$ , see e.g. Kiryakova [18], p. 336,

$$J_\nu^{(\lambda)}(z) := \left(\frac{z}{2}\right)^\nu \sum_{n=0}^{\infty} \frac{(-1)^n (z/2)^{2n}}{n! \Gamma(\lambda n + \nu + 1)} = \left(\frac{z}{2}\right)^\nu W_{\lambda,\nu+1} \left(-\frac{z^2}{4}\right).
 \tag{6}$$

Similar remarks can be extended to the modified Bessel functions  $I_\nu$ .

**Recurrence relations.** Hereafter, we quote some relevant recurrence relations from Erdély et al. [6], Vol. 3, Ch. 18:

$$\begin{aligned}
 \lambda z W_{\lambda,\lambda+\mu}(z) &= W_{\lambda,\mu-1}(z) + (1 - \mu) W_{\lambda,\mu}(z), \\
 \frac{d}{dz} W_{\lambda,\mu}(z) &= W_{\lambda,\lambda+\mu}(z).
 \end{aligned}
 \tag{7}$$

We note that these relations can easily be derived from (1).

### 3. THE AUXILIARY FUNCTIONS $F_\nu(z)$ AND $M_\nu(z)$ IN $\mathbb{C}$

In his earliest analysis of the time fractional diffusion-wave equation [20], Mainardi introduced the two *auxiliary functions* of the Wright type:

$$F_\nu(z) := W_{-\nu,0}(-z) \quad , \quad 0 < \nu < 1,
 \tag{9}$$

and

$$M_\nu(z) := W_{-\nu,1-\nu}(-z) \quad , \quad 0 < \nu < 1,
 \tag{10}$$

inter-related through

$$F_\nu(z) = \nu z M_\nu(z).
 \tag{11}$$

Thus the functions  $F_\nu(z)$  and  $M_\nu$  are particular cases of  $W_{\lambda,\mu}(z)$  by setting  $\lambda = -\nu$  and  $\mu = 0$ ,  $\mu = 1$ , respectively. The motivation was based on the inversion of certain Laplace transforms in order to obtain the fundamental solutions of the fractional diffusion-wave equation in the space-time domain, as shown in Mainardi [20, 21, 22, 23, 24]. Here we will devote particular attention to the mathematical properties of these functions limiting at the essential the discussion for the general

Wright functions. The reader is referred to the Section 7 for some historical and bibliographical details.

**Series representations.** The *series representations* for our auxiliary functions read

$$\begin{aligned}
 (12) \quad F_\nu(z) &:= \sum_{n=1}^{\infty} \frac{(-z)^n}{n! \Gamma(-\nu n)} = \\
 &= \frac{1}{\pi} \sum_{n=1}^{\infty} \frac{(-z)^{n-1}}{n!} \Gamma(\nu n + 1) \sin(\pi \nu n),
 \end{aligned}$$

and

$$\begin{aligned}
 (13) \quad M_\nu(z) &:= \sum_{n=0}^{\infty} \frac{(-z)^n}{n! \Gamma[-\nu n + (1 - \nu)]}, \\
 &:= \frac{1}{\pi} \sum_{n=1}^{\infty} \frac{(-z)^{n-1}}{(n-1)!} \Gamma(\nu n) \sin(\pi \nu n),
 \end{aligned}$$

where we have used the well-known reflection formula for the Gamma function,

$$\Gamma(\zeta) \Gamma(1 - \zeta) = \frac{\pi}{\sin \pi \zeta}.$$

As an exercise the reader can easily prove that the radius of convergence of the power series in (12)-(13) is infinite for  $0 < \nu < 1$ , without be aware of the Wright functions, see also Podlubny [36].

Furthermore we note that  $F_\nu(0) = 0$ ,  $M_\nu(0) = 1/\Gamma(1-\nu)$  and that the relation (11), consistent with the recurrence relation (7), can be derived from (12)-(13) arranging the terms of the series.

**The integral representations.** The *integral representations* for our auxiliary functions read:

$$(14) \quad F_\nu(z) := \frac{1}{2\pi i} \int_{H_a} e^{\sigma - z\sigma^\nu} d\sigma,$$

$$(15) \quad M_\nu(z) := \frac{1}{2\pi i} \int_{H_a} e^{\sigma - z\sigma^\nu} \frac{d\sigma}{\sigma^{1-\nu}}.$$

We note that the relation (11),  $F_\nu(z) = \nu z M_\nu(z)$ , can be obtained directly from (12)-(13) with an integration by parts. In fact,

$$\begin{aligned}
 M_\nu(z) &= \int_{H_a} e^{\sigma - z\sigma^\nu} \frac{d\sigma}{\sigma^{1-\nu}} = \int_{H_a} e^\sigma \left( -\frac{1}{\nu z} \frac{d}{d\sigma} e^{-z\sigma^\nu} \right) d\sigma = \\
 &= \frac{1}{\nu z} \int_{H_a} e^{\sigma - z\sigma^\nu} d\sigma = \frac{F_\nu(z)}{\nu z}.
 \end{aligned}$$

The passage from the series representation to the integral representation and vice-versa for our auxiliary functions can be derived in a way similar to that adopted for the general Wright function, that is by expanding in positive powers of  $z$  the exponential function  $\exp(-z\sigma^\nu)$ , exchanging the order between the series and the integral and using the Hankel representation of the reciprocal of the Gamma function. Since the radius of convergence of the power series in (12)-(13) is infinite for

$0 < \nu < 1$ , our auxiliary functions turn out to be entire in  $z$  and therefore the exchange between the series and the integral is legitimate.

**Special cases.** Explicit expressions of  $F_\nu(z)$  and  $M_\nu(z)$  in terms of known functions are expected for some particular values of  $\nu$ . In Mainardi and Tomirotti [31] the authors have shown that for  $\nu = 1/q$ , where  $q \geq 2$  is a positive integer, the auxiliary functions can be expressed as a sum of  $(q - 1)$  simpler entire functions. In the particular cases  $q = 2$  and  $q = 3$  we find from (13)

$$(16) \quad M_{1/2}(z) = \frac{1}{\sqrt{\pi}} \sum_{m=0}^{\infty} (-1)^m \left(\frac{1}{2}\right)_m \frac{z^{2m}}{(2m)!} = \frac{1}{\sqrt{\pi}} \exp\left(-\frac{z^2}{4}\right),$$

and

$$(17) \quad M_{1/3}(z) = \frac{1}{\Gamma(2/3)} \sum_{m=0}^{\infty} \left(\frac{1}{3}\right)_m \frac{z^{3m}}{(3m)!} - \frac{1}{\Gamma(1/3)} \sum_{m=0}^{\infty} \left(\frac{2}{3}\right)_m \frac{z^{3m+1}}{(3m+1)!} = 3^{2/3} \text{Ai}\left(\frac{z}{3^{1/3}}\right),$$

where Ai denotes the *Airy function*.

Furthermore, it can be proved that  $M_{1/q}(z)$  satisfies the differential equation of order  $q - 1$

$$(18) \quad \frac{d^{q-1}}{dz^{q-1}} M_{1/q}(z) + \frac{(-1)^q}{q} z M_{1/q}(z) = 0,$$

subjected to the  $q - 1$  initial conditions at  $z = 0$ , derived from (15),

$$(19) \quad M_{1/q}^{(h)}(0) = \frac{(-1)^h}{\pi} \Gamma\left(\frac{h+1}{q}\right) \sin \frac{(h+1)\pi}{q},$$

with  $h = 0, 1, \dots, q - 2$ . We note that, for  $q \geq 4$ , equation (18) is akin to the *hyper-Airy* differential equation of order  $q - 1$ , see e.g. Bender and Orszag [2]. Consequently, in view of the above considerations, the auxiliary function  $M_\nu(z)$  can be referred to as the *generalized hyper-Airy function*. However, in view of further applications in stochastic processes, we prefer to consider it as a natural (fractional) generalization of the Gaussian function, see Mura, Taqqu and Manardi [35], Mura and Pagnini [34], Mura and Mainardi [33], similarly as the Mittag-Leffler function is known to be the natural (fractional) generalization of the exponential function for relaxation processes, see Gorenflo and Mainardi [12], Mainardi and Gorenflo [26]. To stress the relevance of the auxiliary function  $M_\nu(z)$ , it was also suggested the special name *M-Wright function*, a terminology that has been followed in literature to some extent, see Notes in Section 7.

#### 4. THE AUXILIARY FUNCTIONS $F_\nu(x)$ AND $M_\nu(x)$ IN $\mathbb{R}$

We point out that the most relevant applications of Wright functions, specially our auxiliary functions, are when the variable is real. More precisely, in this Section, we will consider functions that are defined either on the positive real semi-axis  $\mathbb{R}^+$  or on all of  $\mathbb{R}$ . When the support is  $\mathbb{R}$ , we agree to consider *even functions*, that is, functions defined in a symmetric way. In this case, to stress the symmetry property of the function, the independent variable may be denoted by  $|x|$ .

We point out that in the limit  $\nu \rightarrow 1^-$  the function  $M_\nu(x)$ , for  $x \in \mathbb{R}^+$ , tends to the Dirac generalized function  $\delta(x - 1)$ .

**The asymptotic representation of  $M_\nu(x)$ .** Let us first point out the asymptotic behaviour of the function  $M_\nu(x)$  as  $x \rightarrow +\infty$ . Choosing as a variable  $x/\nu$  rather than  $x$ , the computation of the asymptotic representation by the saddle-point approximation yields, see [31],

$$(20) \quad M_\nu\left(\frac{x}{\nu}\right) \sim a(\nu)x^{(\nu-1/2)/(1-\nu)} \exp\left[-b(\nu)x^{1/(1-\nu)}\right],$$

where

$$(21) \quad a(\nu) = \frac{1}{\sqrt{2\pi(1-\nu)}} > 0, \quad b(\nu) = \frac{1-\nu}{\nu} > 0.$$

The above evaluation is consistent with the first term in Wright's asymptotic expansion (3) after having used the definition (10).

**Plots of  $M_\nu(|x|)$ .** We show the plots of the  $M$ -Wright function on the real axis for some rational values of the parameter  $\nu$ . To gain more insight of the effect of the parameter itself on the behaviour close to and far from the origin, we adopt both linear and logarithmic scale for the ordinates.




FIGURE 1. Plots of the function  $M_\nu(|x|)$  with  $\nu = 0, 1/8, 1/4, 3/8, 1/2$  for  $-5 \leq x \leq 5$ ; left: linear scale, right: logarithmic scale.




FIGURE 2. Plots of the function  $M_\nu(|x|)$  with  $\nu = 1/2, 5/8, 3/4, 1$  for  $-5 \leq x \leq 5$ ; left: linear scale; right: logarithmic scale.

In figures 1 and 2 we compare the plots of the  $M_\nu(x)$  Wright auxiliary functions in  $-5 \leq x \leq 5$  for some rational values in the ranges  $\nu \in [0, 1/2]$  and  $\nu \in [1/2, 1]$ ,

respectively. Thus in figure 1 we see the transition from  $\exp(-|x|)$  for  $\nu = 0$  to  $1/\sqrt{\pi} \exp(-x^2)$  for  $\nu = 1/2$ , whereas in figure 2 we see the transition from  $1/\sqrt{\pi} \exp(-x^2)$  to the delta function  $\delta(1 - |x|)$  for  $\nu = 1$ .

In plotting  $M_\nu(x)$  at fixed  $\nu$  for sufficiently large  $x$  the asymptotic representation (20)-(21) is very useful because, as  $x$  increases, the numerical convergence of the series in (13) becomes poor and poor up to being completely inefficient. Henceforth, the matching between the series and the asymptotic representation is relevant.

### 5. THE LAPLACE TRANSFORM PAIRS

Let us write the Laplace transform of the Wright function as

$$W_{\lambda,\mu}(\pm r) \div \mathcal{L}[W_{\lambda,\mu}(\pm r); s] := \int_0^\infty e^{-sr} W_{\lambda,\mu}(\pm r) dr ,$$

where  $r$  denotes a non negative real variable, i.e.  $0 \leq r < +\infty$ , and  $s$  is the Laplace complex parameter.

When  $\lambda > 0$  the series representation of the Wright function can be transformed term-by-term. In fact, for a known theorem of the theory of the Laplace transforms, see e.g. Doetsch [5], the Laplace transform of an entire function of exponential type can be obtained by transforming term-by-term the Taylor expansion of the original function around the origin. In this case the resulting Laplace transform turns out to be analytic and vanishing at infinity. As a consequence, we obtain *the Laplace transform pair for the Wright function of the first kind* as

$$(22) \quad W_{\lambda,\mu}(\pm r) \div \frac{1}{s} E_{\lambda,\mu} \left( \pm \frac{1}{s} \right) \quad , \quad \lambda > 0 \quad , \quad |s| > 0 \quad ,$$

where  $E_{\lambda,\mu}$  denotes the Mittag-Leffler function in two parameters. The proof is straightforward noting that

$$\sum_{n=0}^\infty \frac{(\pm r)^n}{n! \Gamma(\lambda n + \mu)} \div \frac{1}{s} \sum_{n=0}^\infty \frac{(\pm 1/s)^n}{\Gamma(\lambda n + \mu)} \quad ,$$

and recalling the series representation of the Mittag-Leffler function,

$$E_{\alpha,\beta}(z) := \sum_{n=0}^\infty \frac{z^n}{\Gamma(\alpha n + \beta)} \quad , \quad \alpha > 0 \quad , \quad \beta \in \mathbb{C} \quad .$$

For  $\lambda \rightarrow 0^+$  equation (22) provides the Laplace transform pair

$$(23) \quad W_{0^+,\mu}(\pm r) = \frac{e^{\pm r}}{\Gamma(\mu)} \div \frac{1}{\Gamma(\mu)} \frac{1}{s \mp 1} = \frac{1}{s} E_{0,\mu} \left( \pm \frac{1}{s} \right) \quad , \quad |s| > 1 \quad ,$$

where, to remain in agreement with (22), we have formally put

$$E_{0,\mu}(z) := \sum_{n=0}^\infty \frac{z^n}{\Gamma(\mu)} := \frac{1}{\Gamma(\mu)} E_0(z) := \frac{1}{\Gamma(\mu)} \frac{1}{1-z} \quad , \quad |z| < 1 \quad .$$

We recognize that in this special case the Laplace transform exhibits a simple pole at  $s = \pm 1$  while for  $\lambda > 0$  it exhibits an essential singularity at  $s = 0$ .

For  $-1 < \lambda < 0$  the Wright function turns out to be an entire function of order greater than 1, so that the term-by-term transformation representation is no longer legitimate. Thus, for Wright functions of the second kind, care is required in establishing the existence of the Laplace transform, which necessarily must tend to zero as  $s \rightarrow \infty$  in its half-plane of convergence.

For the sake of convenience we limit ourselves to derive the Laplace transform for the special case of  $M_\nu(r)$ ; the exponential decay as  $r \rightarrow \infty$  of the *original* function provided by (20) ensures the existence of the *image* function. From the integral representation (13) of the  $M_\nu$  function we obtain

$$\begin{aligned} M_\nu(r) &\div \frac{1}{2\pi i} \int_0^\infty e^{-sr} \left[ \int_{Ha} e^{\sigma-r\sigma^\nu} \frac{d\sigma}{\sigma^{1-\nu}} \right] dr = \\ &= \frac{1}{2\pi i} \int_{Ha} e^\sigma \sigma^{\nu-1} \left[ \int_0^\infty e^{-r(s+\sigma^\nu)} dr \right] d\sigma = \frac{1}{2\pi i} \int_{Ha} \frac{e^\sigma \sigma^{\nu-1}}{\sigma^\nu + s} d\sigma . \end{aligned}$$

Then, by recalling the integral representation of the Mittag-Leffler function,

$$E_\alpha(z) = \frac{1}{2\pi i} \int_{Ha} \frac{\zeta^{\alpha-1} e^\zeta}{\zeta^\alpha - z} d\zeta \quad , \quad \alpha > 0 \quad , \quad z \in \mathbb{C} \quad ,$$

we obtain the Laplace transform pair

$$(24) \quad M_\nu(r) := W_{-\nu, 1-\nu}(-r) \div E_\nu(-s) \quad , \quad 0 < \nu < 1 \quad .$$

Although transforming the Taylor series of  $M_\nu(r)$  term-by-term is not legitimate, this procedure yields a series of negative powers of  $s$  that represents the asymptotic expansion of the correct Laplace transform,  $E_\nu(-s)$ , as  $s \rightarrow \infty$  in a sector around the positive real axis. Indeed we get

$$\begin{aligned} \sum_{n=0}^{\infty} \frac{\int_0^\infty e^{-sr} (-r)^n dr}{n! \Gamma(-\nu n + (1-\nu))} &= \sum_{n=0}^{\infty} \frac{(-1)^n}{\Gamma(-\nu n + 1 - \nu)} \frac{1}{s^{n+1}} = \\ &= \sum_{m=1}^{\infty} \frac{(-1)^{m-1}}{\Gamma(-\nu m + 1)} \frac{1}{s^m} \sim E_\nu(-s) \quad , \quad s \rightarrow \infty \quad . \end{aligned}$$

We note that (24) contains the well-known Laplace transform pair, see e.g. Doetsch [5],

$$M_{1/2}(r) := \frac{1}{\sqrt{\pi}} \exp\left(-\frac{r^2}{4}\right) \div E_{1/2}(-s) := \exp s^2 \operatorname{erfc}(s) \quad ,$$

valid  $\forall s \in \mathbb{C}$ .

Analogously, using the more general integral representation (2) of the Wright function, we can get *the Laplace transform pair for the Wright function of the second kind*. For the case  $\lambda = -\nu \in (-1, 0)$ , with  $\mu > 0$  for simplicity, we obtain,

$$(25) \quad W_{-\nu, \mu}(-r) \div E_{\nu, \mu+\nu}(-s) \quad , \quad 0 < \nu < 1 \quad .$$

We note the minus sign in the argument in order to ensure the the existence of the Laplace transform thanks to the Wright asymptotic formula (3) valid in a sector about the negative real axis.

In the limit as  $\nu \rightarrow 0^+$  (thus  $\lambda \rightarrow 0^-$ ) we formally obtain the Laplace transform pair

$$(26) \quad W_{0^-, \mu}(-r) := \frac{e^{-r}}{\Gamma(\mu)} \div \frac{1}{\Gamma(\mu)} \frac{1}{s+1} := E_{0, \mu}(-s) \quad .$$

Therefore, as  $\lambda \rightarrow 0^\pm$ , and  $\mu = 1$  we note a sort of continuity in the results (23) and (26) since

$$(27) \quad W_{0,1}(-r) := e^{-r} \div \frac{1}{s+1} = \begin{cases} \frac{1}{s} E_0(-1/s) & , \quad |s| > 1 ; \\ E_0(-s) & , \quad |s| < 1 . \end{cases}$$

We here point out the relevant *Laplace transform pairs related to the auxiliary functions of argument  $r^{-\nu}$* , see Mainardi [20, 21, 22],

$$(28) \quad \frac{1}{r} F_\nu \left( \frac{1}{r^\nu} \right) = \frac{\nu}{r^{\nu+1}} M_\nu \left( \frac{1}{r^\nu} \right) \div e^{-s^\nu} \quad , \quad 0 < \nu < 1 ,$$

$$(29) \quad \frac{1}{\nu} F_\nu \left( \frac{1}{r^\nu} \right) = \frac{1}{r^\nu} M_\nu \left( \frac{1}{r^\nu} \right) \div \frac{e^{-s^\nu}}{s^{1-\nu}} \quad , \quad 0 < \nu < 1 .$$

We recall that the Laplace transform pairs in (28) were formerly considered by Pollard [37], who provided a rigorous proof based on a formal result by Humbert [15]. Later Mikusiński [32] got a similar result based on his theory of operational calculus, and finally, albeit unaware of the previous results, Buchen and Mainardi [3] derived the result in a formal way. We note, however, that all these authors were not informed about the Wright functions. To our actual knowledge, the former author who derived the Laplace transforms pairs (28)-(29) in terms of Wright functions of the second kind was Stankovič [39].

Hereafter we will provide two independent proofs of (28) by carrying out the inversion of  $\exp(-s^\nu)$ , either by the complex Bromwich integral formula, see Mainardi [20, 21, 22] or by the formal series method, see Buchen and Mainardi [3]. Similarly, we can act for the Laplace transform pair (29).

For the complex integral approach we deform the Bromwich path  $Br$  into the Hankel path  $Ha$ , that is equivalent to the original path, and we set  $\sigma = sr$ . Recalling (13)-(14), we get

$$\begin{aligned} \mathcal{L}^{-1}[\exp(-s^\nu)] &= \frac{1}{2\pi i} \int_{Br} e^{sr-s^\nu} ds = \frac{1}{2\pi i r} \int_{Ha} e^{\sigma-(\sigma/r)^\nu} d\sigma = \\ &= \frac{1}{r} F_\nu \left( \frac{1}{r^\nu} \right) = \frac{\nu}{r^{\nu+1}} M_\nu \left( \frac{1}{r^\nu} \right) . \end{aligned}$$

For the series approach, let us expand the Laplace transform in series of negative powers and invert term by term. Then, after recalling (12)-(13), we obtain:

$$\begin{aligned} \mathcal{L}^{-1}[\exp(-s^\nu)] &= \sum_{n=0}^{\infty} \frac{(-1)^n}{n!} \mathcal{L}^{-1}[s^{\nu n}] = \sum_{\nu=1}^{\infty} \frac{(-1)^n}{n!} \frac{r^{-\nu n-1}}{\Gamma(-\nu n)} = \\ &= \frac{1}{r} F_\nu \left( \frac{1}{r^\nu} \right) = \frac{\nu}{r^{\nu+1}} M_\nu \left( \frac{1}{r^\nu} \right) . \end{aligned}$$

We note the relevance of Laplace transforms (24) and (28) in pointing out the non-negativity of the Wright function  $M_\nu(x)$  and the complete monotonicity of the Mittag-leffler functions  $E_\nu(-x)$  for  $x > 0$  and  $0 < \nu < 1$ . In fact, since  $\exp(-s^\nu)$  denotes the Laplace transform of a probability density (precisely, the extremal Lévy stable density of index  $\nu$ , see Feller [7]), the L.H.S. of (28) must be non-negative, and so also the L.H.S of (24). As a matter of fact the Laplace transform

pair (24) shows, replacing  $s$  by  $x$ , that the spectral representation of the Mittag-Leffler function  $E_\nu(-x)$  is expressed in terms of the  $M$ -Wright function  $M_\nu(r)$ , that is:

$$(30) \quad E_\nu(-x) = \int_0^\infty e^{-rx} M_\nu(r) dr \quad , \quad 0 < \nu < 1 \quad , \quad x \geq 0 .$$

We now recognize that equation (30) is consistent with a result derived by Pollard [37].

It is instructive to compare the spectral representation of  $E_\nu(-x)$  with that of the function  $E_\nu(-t^\nu)$ . We recall

$$(31) \quad E_\nu(-t^\nu) = \int_0^\infty e^{-rt} K_\nu(r) dr \quad , \quad 0 < \nu < 1 \quad , \quad t \geq 0 ,$$

with *spectral function*

$$(32) \quad K_\nu(r) = \frac{1}{\pi} \frac{r^{\nu-1} \sin(\nu\pi)}{r^{2\nu} + 2r^\nu \cos(\nu\pi) + 1} = \frac{1}{\pi r} \frac{\sin(\nu\pi)}{r^\nu + r^{-\nu} + 2 \cos(\nu\pi)} .$$

The relationship between  $M_\nu(r)$  and  $K_\nu(r)$  is worth to be explored. Both functions are non-negative, integrable and normalized in  $\mathbb{R}^+$ , so they can be adopted in probability theory as density functions. The normalization conditions derive from equations (30) and (31) since

$$\int_0^{+\infty} M_\nu(r) dr = \int_0^{+\infty} K_\nu(r) dr = E_\nu(0) = 1 .$$

## 6. THE $M_\nu$ -WRIGHT FUNCTIONS IN PROBABILITY

The  $M_\nu$ -Wright functions play fundamental roles in Theory of Probability and Stochastic Processes with support both in  $\mathbb{R}^+$  (the variable is mostly a time coordinate) and in all of  $\mathbb{R}$  (the function, divided by 2, is continued in a symmetric way so the variable is the absolute value of a space coordinate).

Indeed, for certain stochastic processes of renewal type, functions of Mittag-Leffler and Wright type can be adopted as probability distributions of waiting times, as shown in Mainardi, Gorenflo and Vivoli [27], where such distributions are compared. We refer the interested reader to that paper for details.

As in Section 4, here we agree to denote by  $x$  and  $|x|$  the variable of  $M_\nu$  functions in  $\mathbb{R}^+$  and  $\mathbb{R}$ , respectively.

**The absolute moments of order  $\delta$ .** The *absolute moments* of order  $\delta > -1$  in  $\mathbb{R}^+$  of the Wright  $M$ -function *pdf* in  $\mathbb{R}^+$  are finite and turn out to be

$$(33) \quad \int_0^\infty x^\delta M_\nu(x) dx = \frac{\Gamma(\delta+1)}{\Gamma(\nu\delta+1)} \quad , \quad \delta > -1 \quad , \quad 0 \leq \nu < 1 .$$

In order to derive this fundamental result we proceed as follows, based on the integral representation (15),

$$\begin{aligned} \int_0^\infty x^\delta M_\nu(x) dx &= \int_0^\infty c^\delta \left[ \frac{1}{2\pi i} \int_{Ha} e^{\sigma-x\sigma^\nu} \frac{d\sigma}{\sigma^{1-\nu}} \right] dx = \\ &= \frac{1}{2\pi i} \int_{Ha} e^\sigma \left[ \int_0^\infty e^{-x\sigma^\nu} x^\delta dx \right] \frac{d\sigma}{\sigma^{1-\nu}} = \\ &= \frac{\Gamma(\delta+1)}{2\pi i} \int_{Ha} \frac{e^\sigma}{\sigma^{\nu\delta+1}} d\sigma = \frac{\Gamma(\delta+1)}{\Gamma(\nu\delta+1)}. \end{aligned}$$

Above we have legitimate the exchange between the two integrals and we have used the identity

$$\int_0^\infty e^{-x\sigma^\nu} x^\delta dx = \frac{\Gamma(\delta+1)}{(\sigma^\nu)^{\delta+1}}.$$

In particular, for  $\delta = n \in \mathbb{N}$ , the formula (33) provides the moments of integer order that can also be computed from the Laplace transform pair (24) as follows:

$$\int_0^{+\infty} x^n M_\nu(x) dx = \lim_{s \rightarrow 0} (-1)^n \frac{d^n}{ds^n} E_\nu(-s) = \frac{\Gamma(n+1)}{\Gamma(\nu n+1)}.$$

Incidentally, we note that the Laplace transform pair (24) could be obtained using the fundamental result (33) by developing in power series the exponential kernel of the Laplace transform and then transforming the series term-by-term.

**The characteristic function.** As well-known in probability theory the Fourier transform of a density provides the so-called *characteristic function*. In our case we have:

$$\begin{aligned} \mathcal{F} \left[ \frac{1}{2} M_\nu(|x|) \right] &:= \frac{1}{2} \int_{-\infty}^{+\infty} M_\nu(|x|) dx = \\ (34) \qquad &= \int_0^\infty \cos(\kappa x) M_\nu(x) dx = E_{2\nu}(-\kappa^2). \end{aligned}$$

For this prove it is sufficient to develop in series the cosine function and use formula (33),

$$\begin{aligned} \int_0^\infty \cos(\kappa x) M_\nu(x) dx &= \sum_{n=0}^\infty (-1)^n \frac{\kappa^{2n}}{(2n)!} \int_0^\infty x^{2n} M_\nu(x) dx = \\ &= \sum_{n=0}^\infty (-1)^n \frac{\kappa^{2n}}{\Gamma(2\nu n+1)} = E_{2\nu}(-\kappa^2). \end{aligned}$$

**Relations with Lévy stable distributions.** We find it worth to discuss the relations between the  $M_\nu$ -Wright functions and the so called *Lévy stable distributions*. The term stable has been assigned by the French mathematician Paul Lévy, who in the 1920's years started a systematic research in order to generalize the celebrated *Central Limit Theorem* to probability distributions with infinite variance. For stable distributions we can assume the following

**Definition 1.** If two independent real random variables with the same shape or type of distribution are combined linearly and the distribution of the resulting

random variable has the same shape, the common distribution (or its type, more precisely) is said to be stable.

The restrictive condition of stability enabled Lévy (and then other authors) to derive the *canonic form* for the Fourier transform of the densities of these distributions. Such transform in probability theory is known as *characteristic function*.

Here we follow the parameterization in Feller [7] revisited in Gorenflo and Mainardi [13] and in Mainardi, Luchko and Pagnini [28].

Denoting by  $L_\alpha^\theta(x)$  a generic stable density in  $\mathbb{R}$ , where  $\alpha$  is the *index of stability* and  $\theta$  the asymmetry parameter, improperly called *skewness*, its characteristic function reads:

$$(35) \quad L_\alpha^\theta(x) \div \widehat{L}_\alpha^\theta(\kappa) = \exp[-\psi_\alpha^\theta(\kappa)] \quad , \quad \psi_\alpha^\theta(\kappa) = |\kappa|^\alpha e^{i(\text{sign } \kappa)\theta\pi/2} \quad , \\ 0 < \alpha \leq 2 \quad , \quad |\theta| \leq \min\{\alpha, 2 - \alpha\} \quad .$$

We note that the allowed region for the parameters  $\alpha$  and  $\theta$  turns out to be a diamond in the plane  $\{\alpha, \theta\}$  with vertices in the points  $(0, 0)$ ,  $(1, 1)$ ,  $(1, -1)$ ,  $(2, 0)$ , that we call the *Feller-Takayasu diamond*, see figure 3. For values of  $\theta$  on the border of the diamond (that is  $\theta = \pm\alpha$  if  $0 < \alpha < 1$ , and  $\theta = \pm(2 - \alpha)$  if  $1 < \alpha < 2$ ) we obtain the so-called *extremal stable densities*.

We note the *symmetry relation*  $L_\alpha^\theta(-x) = L_\alpha^{-\theta}(x)$ , so that a stable density with  $\theta = 0$  is symmetric.




FIGURE 3. The Feller-Takayasu diamond for Lévy stable densities.

Stable distributions have noteworthy properties of which the interested reader can be informed from the existing literature. Here-after we recall some peculiar *properties*:

- *The class of stable distributions possesses its own domain of attraction*, see e.g. Feller [7].
- *Any stable density is unimodal and indeed bell-shaped*, i.e. its  $n$ -th derivative has exactly  $n$  zeros in  $\mathbb{R}$ , see Gawronski [8].
- *The stable distributions are self-similar and infinitely divisible*. These properties derive from the canonic form (35) through the scaling property of the Fourier transform.

*Self-similarity* means

$$(36) \quad L_\alpha^\theta(x, t) \div \exp[-t\psi_\alpha^\theta(\kappa)] \iff L_\alpha^\theta(x, t) = t^{-\alpha} L_\alpha^\theta(x/t^\alpha),$$

where  $t$  is a positive parameter. If  $t$  is time, then  $L_\alpha^\theta(x, t)$  is a spatial density evolving on time with self-similarity. *Infinite divisibility* means that for every positive integer  $n$ , the characteristic function can be expressed as the  $n$ th power of some characteristic function, so that any stable distribution can be expressed as the  $n$ -fold convolution of a stable distribution of the same type. Indeed, taking in (35)  $\theta = 0$ , without loss of generality, we have

$$(37) \quad e^{-t|\kappa|^\alpha} = \left[ e^{-(t/n)|\kappa|^\alpha} \right]^n \iff L_\alpha^0(x, t) = [L_\alpha^0(x, t/n)]^{*n},$$

where

$$[L_\alpha^0(x, t/n)]^{*n} := L_\alpha^0(x, t/n) * L_\alpha^0(x, t/n) * \dots * L_\alpha^0(x, t/n)$$

is the multiple Fourier convolution in  $\mathbb{R}$  with  $n$  identical terms.

Only for a few particular cases, the inversion of the Fourier transform in (35) can be carried out using standard tables, and well-known probability distributions are obtained.

For  $\alpha = 2$  (so  $\theta = 0$ ), we recover the *Gaussian pdf*, that turns out to be the only stable density with finite variance, and more generally with finite moments of any order  $\delta \geq 0$ . In fact

$$(38) \quad L_2^0(x) = \frac{1}{2\sqrt{\pi}} e^{-x^2/4}.$$

All the other stable densities have finite absolute moments of order  $\delta \in [-1, \alpha)$  as we will show later.

For  $\alpha = 1$  and  $|\theta| < 1$ , we get

$$(39) \quad L_1^\theta(x) = \frac{1}{\pi} \frac{\cos(\theta\pi/2)}{[x + \sin(\theta\pi/2)]^2 + [\cos(\theta\pi/2)]^2},$$

which for  $\theta = 0$  includes the *Cauchy-Lorentz pdf*,

$$(40) \quad L_1^0(x) = \frac{1}{\pi} \frac{1}{1+x^2}.$$

In the limiting cases  $\theta = \pm 1$  for  $\alpha = 1$  we obtain the *singular Dirac pdf's*

$$(41) \quad L_1^{\pm 1}(x) = \delta(x \pm 1).$$

In general, we must recall the power series expansions provided in Feller [7]. We restrict our attention to  $x > 0$  since the evaluations for  $x < 0$  can be obtained using the symmetry relation. The convergent expansions of  $L_\alpha^\theta(x)$  ( $x > 0$ ) turn out to be

for  $0 < \alpha < 1$ ,  $|\theta| \leq \alpha$ :

$$(42) \quad L_\alpha^\theta(x) = \frac{1}{\pi x} \sum_{n=1}^{\infty} (-x^{-\alpha})^n \frac{\Gamma(1+n\alpha)}{n!} \sin\left[\frac{n\pi}{2}(\theta-\alpha)\right];$$

for  $1 < \alpha \leq 2$ ,  $|\theta| \leq 2-\alpha$ :

$$(43) \quad L_\alpha^\theta(x) = \frac{1}{\pi x} \sum_{n=1}^{\infty} (-x)^n \frac{\Gamma(1+n/\alpha)}{n!} \sin\left[\frac{n\pi}{2\alpha}(\theta-\alpha)\right].$$

From the series in (42) and the symmetry relation we note that *the extremal stable densities for  $0 < \alpha < 1$  are unilateral*, precisely vanishing for  $x > 0$  if  $\theta = \alpha$ , vanishing for  $x < 0$  if  $\theta = -\alpha$ . In particular the unilateral extremal densities  $L_{\alpha}^{-\alpha}(x)$  with  $0 < \alpha < 1$  have support in  $\mathbb{R}^+$  and Laplace transform  $\exp(-s^{\alpha})$ . For  $\alpha = 1/2$  we obtain the so-called *Lévy-Smirnov pdf*:

$$(44) \quad L_{1/2}^{-1/2}(x) = \frac{x^{-3/2}}{2\sqrt{\pi}} e^{-1/(4x)} \quad , \quad x \geq 0 .$$

As a consequence of the convergence of the series in (42)-(43) and of the symmetry relation we recognize that the stable *pdf*'s with  $1 < \alpha \leq 2$  are entire functions, whereas with  $0 < \alpha < 1$  have the form

$$(45) \quad L_{\alpha}^{\theta}(x) = \begin{cases} \frac{1}{x} \Phi_1(x^{-\alpha}) & \text{for } x > 0 , \\ \frac{1}{|x|} \Phi_2(|x|^{-\alpha}) & \text{for } x < 0 , \end{cases}$$

where  $\Phi_1(z)$  and  $\Phi_2(z)$  are distinct entire functions. The case  $\alpha = 1$  ( $|\theta| < 1$ ) must be considered in the limit for  $\alpha \rightarrow 1$  of (42)-(43), because the corresponding series reduce to power series akin with geometric series in  $1/x$  and  $x$ , respectively, with a finite radius of convergence. The corresponding stable *pdf*'s are no longer represented by entire functions, as can be noted directly from their explicit expressions (39)-(40).

We omit to provide the asymptotic representations of the stable densities referring the interested reader to Feller [7], Mainardi, Luchko and Pagnini [28]. However, based on asymptotic representations, we can state as follows; for  $0 < \alpha < 2$  the stable *pdf*'s exhibit *fat tails* in such a way that their absolute moment of order  $\delta$  is finite only if  $-1 < \delta < \alpha$ . More precisely, one can show that for non-Gaussian, not extremal, stable densities the asymptotic decay of the tails is

$$(46) \quad L_{\alpha}^{\theta}(x) = O\left(|x|^{-(\alpha+1)}\right) \quad , \quad x \rightarrow \pm\infty .$$

For the extremal densities with  $\alpha \neq 1$  this is valid only for one tail (as  $|x| \rightarrow \infty$ ), the other (as  $|x| \rightarrow \infty$ ) being of exponential order. For  $1 < \alpha < 2$  the extremal *pdf*'s are two-sided and exhibit an exponential left tail (as  $x \rightarrow -\infty$ ) if  $\theta = +(2-\alpha)$ , or an exponential right tail (as  $x \rightarrow +\infty$ ) if  $\theta = -(2-\alpha)$ . Consequently, the Gaussian *pdf* is the unique stable density with finite variance. Furthermore, when  $0 < \alpha \leq 1$ , the first absolute moment is infinite so we should use the median instead of the non-existent expected value in order to characterize the corresponding *pdf*.

Let us also recall a relevant identity between stable densities with index  $\alpha$  and  $1/\alpha$  (a sort of reciprocity relation) pointed out in Feller [7], that is, assuming  $x > 0$ ,

$$(47) \quad \frac{1}{x^{\alpha+1}} L_{1/\alpha}^{\theta}(x^{-\alpha}) = L_{\alpha}^{\theta^*}(x) \quad , \quad \frac{1}{2} \leq \alpha \leq 1 \quad , \quad \theta^* = \alpha(\theta + 1) - 1 .$$

The condition  $1/2 \leq \alpha \leq 1$  implies  $1 \leq 1/\alpha \leq 2$ . A check shows that  $\theta^*$  falls within the prescribed range  $|\theta^*| \leq \alpha$  if  $|\theta| \leq 2 - 1/\alpha$ . We leave as an exercise for the interested reader the verification of this reciprocity relation in the limiting cases  $\alpha = 1/2$  and  $\alpha = 1$ .

From a comparison between the series expansions in (42)-(43) and in (14)-(15), we recognize that for  $x > 0$  our *auxiliary functions of the Wright type are related to*

the extremal stable densities as follows, see [31],

$$(48) \quad L_\alpha^{-\alpha}(x) = \frac{1}{x} F_\alpha(x^{-\alpha}) = \frac{\alpha}{x^{\alpha+1}} M_\alpha(x^{-\alpha}) \quad , \quad 0 < \alpha < 1 \quad ,$$

$$(49) \quad L_\alpha^{\alpha-2}(x) = \frac{1}{x} F_{1/\alpha}(x) = \frac{1}{\alpha} M_{1/\alpha}(x) \quad , \quad 1 < \alpha \leq 2 \quad .$$

In equations (48)-(49), for  $\alpha = 1$ , the skewness parameter turns out to be  $\theta = -1$ , so we get the singular limit

$$(50) \quad L_1^{-1}(x) = M_1(x) = \delta(x - 1) \quad .$$

More generally, all (regular) stable densities, given in equations (42)-(43), were recognized to belong to the class of Fox  $H$ -functions, as formerly shown by Schneider [38]. This general class of high transcendental functions is out of the scope of this survey.

**The Wright  $\mathbb{M}$ -function in two variables.** In view of time-fractional diffusion processes related to time-fractional diffusion equations it is worthwhile to introduce the function in two variables

$$(51) \quad \mathbb{M}_\nu(x, t) := t^{-\nu} M_\nu(xt^{-\nu}) \quad , \quad 0 < \nu < 1 \quad , \quad x, t \in \mathbb{R}^+ \quad ,$$

which defines a spatial probability density in  $x$  evolving in time  $t$  with self-similarity exponent  $H = \nu$ . Of course for  $x \in \mathbb{R}$  we have to consider the symmetric version obtained from (51) multiplying by  $1/2$  and replacing  $x$  by  $|x|$ .

Hereafter we provide a list of the main properties of this function, which can be derived from the Laplace and Fourier transforms for the corresponding Wright  $M$ -function in one variable.

From equation (29) we derive the Laplace transform of  $\mathbb{M}_\nu(x, t)$  with respect to  $t \in \mathbb{R}^+$ ,

$$(52) \quad \mathcal{L} \{ \mathbb{M}_\nu(x, t); t \rightarrow s \} = s^{\nu-1} e^{-xs^\nu} \quad .$$

From equation (24) we derive the Laplace transform of  $\mathbb{M}_\nu(x, t)$  with respect to  $x \in \mathbb{R}^+$ ,

$$(53) \quad \mathcal{L} \{ \mathbb{M}_\nu(x, t); x \rightarrow s \} = E_\nu(-st^\nu) \quad .$$

From equation (34) we derive the Fourier transform of  $\mathbb{M}_\nu(|x|, t)$  with respect to  $x \in \mathbb{R}$ ,

$$(54) \quad \mathcal{F} \{ \mathbb{M}_\nu(|x|, t); x \rightarrow \kappa \} = 2E_{2\nu}(-\kappa^2 t^\nu) \quad .$$

Using the Mellin transforms, Mainardi, Pagnini and Gorenflo [30] the following integral formula,

$$(55) \quad \mathbb{M}_\nu(x, t) = \int_0^\infty \mathbb{M}_\lambda(x, \tau) \mathbb{M}_\mu(\tau, t) d\tau \quad , \quad \nu = \lambda\mu \quad .$$

Special cases of the Wright  $\mathbb{M}$ -function are simply derived for  $\nu = 1/2$  and  $\nu = 1/3$  from the corresponding ones in the complex domain, see equations (16)-(17). We devote particular attention to the case  $\nu = 1/2$  for which we get from (16) the Gaussian density in  $\mathbb{R}$ ,

$$(56) \quad \mathbb{M}_{1/2}(|x|, t) = \frac{1}{2\sqrt{\pi}t^{1/2}} e^{-x^2/(4t)} \quad .$$

For the limiting case  $\nu = 1$  we obtain

$$(57) \quad \mathbb{M}_1(|x|, t) = \frac{1}{2} [\delta(x - t) + \delta(x + t)] .$$

## 7. NOTES

In early nineties, in his former analysis of fractional equations interpolating diffusion and wave-propagation, Mainardi [20] introduced the functions of the Wright type

$$F_\nu(z) := W_{-\nu,0}(-z) \quad , \quad M_\nu(z) := W_{-\nu,1-\nu}(-z) ,$$

with  $0 < \nu < 1$ , inter-related through  $F_\nu(z) = \nu z M_\nu(z)$  to characterize the solutions for typical boundary value problems.

Being in that time only aware of the Bateman project where the parameter  $\lambda$  of the Wright function  $W_{\lambda,\mu}(z)$  was erroneously restricted to non-negative values, Mainardi thought to have extended the original Wright function, in an original way, calling  $F_\nu$  and  $M_\nu$  *auxiliary functions*. Presumably for this reason the function  $M_\nu$  is referred as the *Mainardi function* in the book Podlubny [36] and in some papers including Balescu [1], Chechkin et al. [4], Germano et al. [9], Gorenflo, Luchko and Mainardi [10, 11], Hanyga [14].

It was Professor Stanković, during the presentation of the paper Mainardi and Tomirotti [31] at the Conference *Transform Methods and Special Functions, Sofia 1994*, who informed Mainardi that this extension for  $-1 < \mu < 0$  was already made just by Wright himself in 1940 (following his previous papers in 1930's). In a paper devoted to the 80-th birthday of Prof. Stanković, see Mainardi, Gorenflo and Vivoli [27], Mainardi took the occasion to renew his personal gratitude to Prof. Stanković for this earlier information that has induced him to study the original papers by Wright and work (also in collaboration) on the functions of the Wright type for further applications, see e.g. Gorenflo, Luchko and Mainardi [10, 11] and Mainardi and Pagnini [29].

For more mathematical details on the functions of the Wright type, the reader may be referred to the article by Kilbas, Saigo and Trujillo [16] and to the book Kilbas, Srivastava and Trujillo [17] and references therein. For the numerical point of view we like to point out the recent paper by Luchko [19], where algorithms are provided for computation of the Wright function on the real axis with prescribed accuracy.

## REFERENCES

- [1] R. Balescu, *V-Langevin equations, continuous time random walks and fractional diffusion*, Chaos, Solitons and Fractals, 34(2007), 62–80.
- [2] C.M. Bender & S.A. Orszag, *Advanced mathematical methods for scientists and engineers*, McGraw-Hill, Singapore, 1987.
- [3] P.W. Buchen, & F. Mainardi, *Asymptotic expansions for transient viscoelastic waves*, Journal de Mécanique, 4(1975), 597–608.
- [4] A.V. Chechkin, V.Y. Gonchar, R. Gorenflo, M. Korabel & I.M. Sokolov, *Generalized fractional diffusion equations for accelerating subdiffusion and truncated Lévy flights*, Phys. Rev. E, 78(2008), 021111, 1–13.
- [5] G. Doetsch, *Introduction to the theory and application of the Laplace transformation*, Springer Verlag, Berlin, 1974.
- [6] A. Erdélyi, W. Magnus, F. Oberhettinger & F.G. Tricomi, *Higher transcendental functions*, Vol. 3, McGraw-Hill, New York, 1953-1955.
- [7] W. Feller, *An introduction to probability theory and its applications*, Wiley, New York, Vol. 2, Second Edition, 1971.
- [8] W. Gawronski, *On the bell-shape of stable densities*, Ann. Probab., 12(1984), 230–242.

- [9] G. Germano, M. Politi, E. Scalas & R.E. Schilling, *Stochastic calculus for uncoupled continuous-time random walks*, Phys. Rev. E, 79(2009), 066102, 1–12.
- [10] R. Gorenflo, Y. Luchko & F. Mainardi, *Analytical properties and applications of the Wright function*, Fract. Calc. Appl. Anal., 2(1999), 383–414.
- [11] R. Gorenflo, Y. Luchko & F. Mainardi, *Wright functions as scale-invariant solutions of the diffusion-wave equation*, J. Comput. Appl. Math., 118(2000), 175–191.
- [12] R. Gorenflo & F. Mainardi, *Fractional calculus: integral and differential equations of fractional order*, in *Fractals and fractional calculus in continuum mechanics*, A. Carpinteri & F. Mainardi, (Ed.), Springer Verlag, Wien, 1997, 223–276; <http://arxiv.org/abs/0805.3823>.
- [13] R. Gorenflo & F. Mainardi, *Random walk models for space-fractional diffusion processes*, Fract. Calc. Appl. Anal., 1(1998), 167–191.
- [14] A. Hanyga, *Multidimensional solutions of time-fractional diffusion-wave equations*, Proc. R. Soc. London A, 458(2002), 933–957.
- [15] P. Humbert, *Nouvelles correspondances symboliques*, Bull. Sci. Math. (2), 69(1945), 121–129.
- [16] A.A. Kilbas, M. Saigo & J.J. Trujillo, *On the generalized Wright function*, Fract. Calc. Appl. Anal., 5(4)(2002), 437–460.
- [17] A.A. Kilbas, H.M. Srivastava & J.J. Trujillo, *Theory and applications of fractional differential equations*, North-Holland Series on Mathematics Studies, vol. 204, Elsevier, Amsterdam, 2006.
- [18] V. Kiryakova, *Generalized fractional calculus and applications*, Pitman Research Notes in Mathematics, vol. 301, Longman, Harlow, 1994.
- [19] Y. Luchko, *Algorithms for evaluation of the Wright function for the real arguments' values*, Fract. Calc. Appl. Anal., 11(2008), 57–75.
- [20] F. Mainardi, *On the initial value problem for the fractional diffusion-wave equation*, in *Waves and stability in continuous media*, S. Rionero & T. Ruggeri, (Ed.) , World Scientific, Singapore, 1994, 246–251.
- [21] F. Mainardi, *The fundamental solutions for the fractional diffusion-wave equation*, Appl. Math. Lett., 9(6)1996, 23–28.
- [22] F. Mainardi, *Fractional relaxation-oscillation and fractional diffusion-wave phenomena*, Chaos, Solitons & Fractals, 7(1996), 1461–1477.
- [23] F. Mainardi, *Fractional calculus: some basic problems in continuum and statistical mechanics*, in *Fractals and fractional calculus in continuum mechanics*, A. Carpinteri & F. Mainardi (Ed.) , Springer Verlag, Wien, 1997, 291–348; <http://www.fractalmo.org>.
- [24] F. Mainardi, *Fractional calculus and waves in linear viscoelasticity*, Imperial College Press, London, 2010; <http://www.icpress.co.uk/mathematics/p614.html>.
- [25] F. Mainardi & R. Gorenflo, *On Mittag-Leffler-type functions in fractional evolution processes*, J. Comput. Appl. Math., 118(2000), 283–299.
- [26] F. Mainardi & R. Gorenflo, *Time-fractional derivatives in relaxation processes: a tutorial survey*, Fract. Calc. Appl. Anal., 10(2007), 269–308.
- [27] F. Mainardi, R. Gorenflo & A. Vivoli, *Renewal processes of Mittag-Leffler and Wright type*, Fract. Calc. Appl. Anal., 8(2005), 7–38; <http://arxiv.org/abs/math/0701455>.
- [28] F. Mainardi, Y. Luchko & G. Pagnini, *The fundamental solution of the space-time fractional diffusion equation*, Fract. Calc. Appl. Anal., 4(2001), 153–192; <http://arxiv.org/abs/cond-mat/0702419>.
- [29] F. Mainardi & G. Pagnini, *The Wright functions as solutions of the time-fractional diffusion equations*, Appl. Math. Comp., 141(2003), 51–62.
- [30] F. Mainardi, G. Pagnini & R. Gorenflo, *Mellin transform and subordination laws in fractional diffusion processes*, Fract. Calc. and Appl. Anal., 6(4)(2003), 441–459; <http://arxiv.org/abs/math/0702133>.
- [31] F. Mainardi & M. Tomirotti, *On special function arising in the time fractional diffusion-wave equation*, in *Transform methods and special functions*, Sofia, 1994, P. Rusev, I. Dimovski & V. Kiryakova (Ed.), Science Culture Technology, Singapore, 1995, 171–183.
- [32] J. Mikusiński, *On the function whose Laplace transform is  $\exp(-s^\alpha)$* , Studia Math., 18(1959), 191–198.

- [33] A. Mura & F. Mainardi, *A class of self-similar stochastic processes with stationary increments to model anomalous diffusion in physics*, Integral Transforms Spec. Funct., 20(3-4)(2009), 185–198; <http://arxiv.org/abs/0711.0665>.
- [34] A. Mura & G. Pagnini, *Characterizations and simulations of a class of stochastic processes to model anomalous diffusion*, J. Phys. A: Math. Theor., 41(2008), 285003, 1–22; <http://arxiv.org/abs/0801.4879>.
- [35] A. Mura, M.S. Taqqu & F. Mainardi, *Non-Markovian diffusion equations and processes: analysis and simulation*, Physica A, 387(2008), 5033–5064; <http://arxiv.org/abs/0712.0240>.
- [36] I. Podlubny, *Fractional differential equations*, Mathematics in Science and Engineering, vol. 198, Academic Press, San Diego, 1999.
- [37] H. Pollard, *The representation of  $\exp(-x^\lambda)$  as a Laplace integral*, Bull. Amer. Math. Soc., 52(1946), 908–910.
- [38] W.R. Schneider, *Stable distributions: Fox function representation and generalization*, in *Stochastic processes in classical and quantum systems*, S. Albeverio, G. Casati & D. Merlini, (Ed.), Lecture Notes in Physics, vol. 262, Springer Verlag, Berlin, 1986, 497–511.
- [39] B. Stanković, *On the function of E.M. Wright*, Publ. de l'Institut Mathématique, Beograd, Nouvelle Sér., 10(1970), 113–124.
- [40] R. Wong Y.-Q. Zhao, *Smoothing of Stokes' discontinuity for the generalized Bessel function*, Proc. R. Soc. London A, 455(1999), 1381–1400.
- [41] R. Wong & Y.-Q. Zhao, *Smoothing of Stokes' discontinuity for the generalized Bessel function II*, Proc. R. Soc. London A, 455(1999), 3065–3084.
- [42] E.M. Wright, *On the coefficients of power series having exponential singularities*, Journal London Math. Soc., 8(1933), 71–79.
- [43] E.M. Wright, *The asymptotic expansion of the generalized Bessel function*, Proc. London Math. Soc. (Ser. II), 38(1935), 257–270.
- [44] E.M. Wright, *The asymptotic expansion of the generalized hypergeometric function*, Journal London Math. Soc., 10(1935), 287–293.
- [45] E.M. Wright, *The generalized Bessel function of order greater than one*, Quart. J. Math., Oxford Ser., 11(1940), 36–48.